

*The Active Adult with  
OSTEOARTHRITIS*

Karl B. Fields, MD  
Professor of Family Medicine and Sport Medicine UNC  
Cone Health Sports Medicine Fellowship  
Greensboro AHEC  
2020

---

---

---

---

---

---

---

---

1

*Osteoarthritis*

The most common arthritis  
worldwide affecting a majority of  
persons 65 and over

---

---

---

---

---

---

---

---

2

*Common symptoms of OA*

- Stiffness and decreased joint motion
- Pain – progresses through stages
- Swelling
- Joint deformity
- Instability

---

---

---

---

---

---

---

---

3

### *Generalized OA*

- Involvement of hands with Heberden and Bouchard nodes
  - Multiple Heberden nodes is key marker
  - Onset usually in middle age
- Involvement of spinal joints
- Involvement of 2 other joints
- Not likely a specific syndrome so much as a genetic susceptibility to OA

---

---

---

---

---

---

---

---

4

### *Osteoarthritis Changes*

- Process of continual destruction and repair
- Subchondral bone becomes thickened, sclerotic and eburnation occurs - pain from repetitive microfractures
- Subchondral cysts
- Osteophytes

---

---

---

---

---

---

---

---

5

### *Locations of Osteoarthritis*

- Hip
- Knee
- Cervical and lumbar spine
- CMC, DIP, PIP and first MCP joint of hands
- Foot and Ankle

---

---

---

---


---

---

---

---

6



7

---

---

---

---

---

---

---

---



8

---

---

---

---

---

---

---

---



9

---

---

---

---

---

---

---

---



10

---

---

---


---

---

---

---

---



11

---

---

---


---

---

---

---

---



12

---

---

---


---

---

---

---

---



13

---

---

---

---

---

---

---

---



14

---

---

---

---

---

---

---

---

*Diagnosis of OA*

- Physical exam and XR are typically adequate
- MRI, CT and US useful in specific cases
- US and MR have shown that CPPD may complicate 30% of cases increasing to 60% in older patients
- CPPD crystals can create more inflammatory and pain issues

15

---

---

---

---

---

---

---

---

### *Treating the Patient with OA*

- Rx is oriented toward maintaining function
- Reduction of pain
- Limiting ongoing injury
- Decreasing risk of complications such as surgical intervention or permanent joint damage

---

---

---

---

---

---

---

---

16

### *Nonpharmacologic Approaches to RX*

- Emphasize exercise to maintain aerobic fitness
- Decrease stress of ADLs
- Weight loss when indicated
- Range of motion
- Improve strength

---

---

---

---

---

---

---

---

17

### *Changing Paradigm of OA and Exercise*

- Old Paradigm: joint is damaged so maximal rest and limited stress is best strategy
- New Paradigm is POLICE
  - Protection
  - Optimal Loading
  - Ice
  - Compression
  - Elevation

---

---

---

---

---

---

---

---

18

### *Exercise as Therapy for OA*

- Beneficial effect in OA of knee documented in 17 studies to reduce pain and improve function.
- Guidelines for optimal exercise type and dose are lacking but both biking and water exercise were very effective in Knee OA
- Trend to allow a mix of weight bearing – walking or hiking, strength work, Pilates or yoga, biking or swimming
- Alternate weight bearing and non WB days

---

---

---

---

---

---

---

---

19

### *Does Exercise Cause OA?*

- Studies of runners suggest that risk is actually decreased
- Studies linking exercise to OA usually show that injury was actually the key factor
- Occupational studies probably demonstrate the same confounder

---

---

---

---

---

---

---

---

20

### *Jama 2020: Risks for Knee OA*

- Large population study looked at activity levels and knee OA – 10 year f/u Xrays
- All subjects had mild symptoms and high function
- Low to moderate physical activity. OR 0.69
- Any vigorous activity OR 0.75
- Long term extensive sitting OR 1.0

---

---

---

---

---

---

---

---

21

## *Jama 2020: physical activity and knee OA*

- 49.7% failed to perform any strenuous P.A. in 8 years
- 42.5% reported persistent moderate-to-high frequency of extensive sitting.
- Specific factors limit physical activity
  - Older age
  - higher BMI
  - more severe knee pain
  - non-college-graduate education level
  - weaker quadriceps
  - depression

22

---

---

---

---

---

---

---

---

## *OARSI guidelines for Medical treatment*

- Topical non-steroidal NSAIDs were strongly recommended for individuals with Knee OA (Level 1A).
- OA and GI symptoms: COX-2 inhibitors were Level 1B and NSAIDs with proton pump inhibitors Level 2.
- OA and CVD or frailty: oral NSAIDs not recommended. Intra-articular (IA) corticosteroids, IA
- OA and CVD: hyaluronic acid, and aquatic exercise were Level 1B/Level 2 treatments for Knee OA, but not hip or other areas.

23

---

---

---

---

---

---

---

---

## *NSAIDS*

- Used at both analgesic and therapeutic doses for OA – trend is for use minimum amount required
- Synovitis is rare so most patients do not require therapeutic dosing or continuous therapy
- Patients respond differently to various classes so change of drug should switch class
- Major drawback is the GI toxicity of all products and risk in patients with CVD or renal disease

24

---

---

---

---

---

---

---

---



### *Placebo, Tylenol, Opioids*

- Acetaminophen/Paracetamol (APAP) was conditionally not recommended (Level 4A and 4B),
- use of oral and transdermal opioids was strongly not recommended
- Placebo response benefited 60% of patients

25

---

---

---

---

---

---

---

---

### *Duloxetine*

- Pain relief from central nociceptive pathways by selective inhibition of serotonin and norepinephrine reuptake
- Pain relief demonstrated in RCTs with RR 1.49 and 1.69 of 30 to 50% pain reduction
- Start with 30 mg but effective at 60 to 120 mg per day
- Nausea in 15 %, fatigue , dizziness, dry mouth

26

---

---

---

---

---

---

---

---

### *Tramadol for OA*

- Downgraded benefit for pain to questionable clinical benefit *Cochrane 2019*
- Up to 50% more patients do experience a 20% or greater improvement than placebo
- Still only 20% of patients have strong response
- Side-effects led to RR of 2.6 of drop out

27

---

---

---

---

---

---

---

---

### *Corticosteroid Injection*

- Studies suggest pain relief from 1 to 6 weeks with decline to no benefit by 13 weeks.
- NNT 8 with either 40 or 80 of triamcinolone or methylprednisolone
- By 6 months no difference shown in placebo-controlled trials
- Risk and side-effects minimal
- Long term PT and exercise programs provided as much pain relief

---

---

---

---

---

---

---

---

28

### *Hyaluranon – No longer recommended by AAOS*

- Hyaluranon showed minimal benefit for pain vs. placebo that did not clearly meet clinically important difference
- Hyaluranon vs. CSI no clear benefit
- Side-effects of flares and wallet damage

---

---

---

---

---

---

---

---

29

### *Platelet Rich Plasma*

- Meta-analysis of 10 trials showed significant benefit in pain reduction
- PRP outperformed placebo
- PRP outperformed Hyaluranon
- Trials had high risk of bias and unclear how many injections needed
- No long term or outcome-based EBM yet
- Relatively expensive

---

---

---

---

---

---

---

---

30

## Supplements – limited EBM

- Most supplements show a lack of clinically important benefit
  - glucosamine, chondroitin,
  - vitamin D, diacerein, avocado soybean unsaponifiables (ASU),
  - fish oil
- curcumin (active ingredient of turmeric) small trials show benefit and no significant side-effects
- Boswellia serrata small trials with benefit and no significant side-effects

31

---

---

---

---

---

---

---

---

## Activity After Knee Arthroplasty

- Diduch: 88 pts/ 114 total knees/ mean age 51
- 86 improved activity/ 24% to vigorous sport
- Bradbury: 159 pts/ 208 knees
- 65% returned to sports activity
- Bowling 91%/ golf 57%/ tennis 20%



32

---

---

---

---

---

---

---

---

## Summary OA

- OA affects a majority of people sometime in life
- Exercise, weight loss and overall conditioning provide the cornerstone for successful disease control
- Medical options offer relief of pain
- Tylenol and Hyaluranon have lost favor
- TKR and THR returns older people to sport

33

---

---

---

---

---

---

---

---